

**PRECAUTIONS:** Avoid inhalation and skin contact. Wash contaminated skin and clothes. Destroy empty containers after use. Keep away from children, animal, feed and food items.

**PROTECTIVE GEAR:** When mixing or applying, wear rubber gloves, goggles, respirator, boots, socks and protective clothing. Remove protective clothing and wash hands and face thoroughly before meals and after work.

**Physical & Chemical Hazards:** Sensitive to heat. Avoid contact with highly reactive chemicals such as strong acids, strong base or strong oxidising agent.

**Symptoms of Poisoning:** May cause mild to moderate irritation to skin and eyes. May be irritating to the respiratory system. If ingested may cause nausea, vomiting and intestinal spasms.

**STORAGE AND DISPOSAL:** Store in a cool, dry, well ventilated place. Do not use or store near heat, open flames or hot surfaces. Store in original containers only. Keep out of the reach of the children and animals. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal. Triple rinse then crush or puncture container and dispose of in a sanitary landfill or incinerator. DO NOT re-use product containers. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and setup for this purpose, clear off waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### FIRST AID

**Eyes:** Hold the eyes open and flush with a steady, gentle stream of clean water for 15 minutes. Seek medical attention.

**Skin:** Take off contaminated protective clothing wash the skin thoroughly with soap and clean water. Seek medical attention.

**Ingestion:** Do not induce vomiting. Do not give anything to unconscious patient. Wash (rinse) the mouth and keep the patient calm. Seek medical attention.

**Inhalation:** Remove exposed person to fresh air. Seek medical attention.

**Antidote:** No specific antidote, treat symptomatically.

**Note to Physician:** Treat symptomatically also contact Caribbean Poison Information Network (CARPIN): 1 888 764 7667 or (876) 927-1680 ext. 2300 or 618-1152



[Soluble Liquid]

Systemic Fungicide

#### Composition:

Active Ingredient:	w/v
Metalaxyl	7.00%
Inert Ingredient	93.00%
Total	100.00%



KEEP OUT OF REACH OF CHILDREN  
READ LABEL AND LEAFLET BEFORE USE

Distributed by



NEWPORT-FERSAN (JAMAICA) LTD

2a/2b Wherry Wharf Complex  
Kingston 15, Jamaica W.I.

Tel: 876-967-5815 | Fax: 876 -948-9078

Manufactured by

PARIJAT INDUSTRIES PVT. LTD

M-77, Block Market, Greater Kailash - II,

New Delhi 110048, India

**RECOMMENDED USES:** VICTORY + is a systemic fungicide. The mechanism of action involves inhibition of protein synthesis in fungi, by interference with the synthesis of ribosomal RNA. Fungal population infesting the host is controlled by protective and curative action of the fungicide. The systemic action provides complete translocation of fungicide in the plant to ensure effective control of all diseased plant parts. Re-infestation by fungus is limited by protective action of fungicide.

**Environmental Hazard:** Very toxic to fish, hence, harmful to aquatic organism, therefore, not to be used near aquaculture and Rice- Fish farming. Do not contaminate lakes, rivers, ponds and streams.

**Preparation of spray solution:** Mix the recommended dose of VICTORY + in ¼ of the recommended quantity of water to the spray tank while agitating. Add remaining quantity of water with continuous agitation. Ensure complete dispersion of the product in mix water before spraying of the spray solution. Maintain agitation while spraying.

**Time of application:** When weather conditions favour disease development but before infection occur. The crop should be re-sprayed at specified intervals.

**Spray interval:** 10-14 days

**Re-entry interval:** Minimum of 48 hours for re-entry period is necessary to avoid occupational hazard.

**Upon purchase or use of this product, purchaser and user agree to terms of warranty.**

**WARRANTY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer, and the limit of the liability of this company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product (including claims based in contract, negligence, strict liability, other tort or otherwise) shall be the purchase price paid by the user or buyer for the quantity of this product involved, or, at the election of this company or any other seller, the replacement of such quantity, or, if not acquired by purchase, replacement of such quantity. In no event shall this company or any other seller be liable for any incidental, consequential or special damages.

#### APPLICATION RATES

Crops/Pests		Application Rate	Preharvest Interval (PHI)	Additional information
Cucumbers, Watermelons, Melons, Squash, Pumpkin, Canteloupe	Downey Mildew ( <i>Pseudoperonospora spp.</i> )	3.0L-4.0L/Ha 1.2L-2L/acre 120ml-200ml/square (Use 500L of water/Ha)	8 days	Begin spraying when weather conditions favour downy mildew development, but before infection has occurred. Apply max. 4 sprays per season. Follow the 10 day interval during rainy periods with heavy disease pressure.
Lettuce, Onions (green), Sweet Potato			20 days	
Cabbage, Cauliflower, Broccoli, Brussel Sprouts, Kale			8 days	
Turnip, Mustard, Collards			20 days	
Tomato	Early Blight ( <i>Alternaria solani</i> )	3.0L-4.0L/Ha 1.2L-2L/acre 120ml-200ml/square (Use 500L of water/Ha)	5 days	Use for the first 4 sprays during the period of active growth. Protective application; 7-10 day interval in situation With heavy disease pressure.
Irish Potato (Solonaceae)	Late Blight ( <i>Phytophthora infestans</i> )		20 days	
(Pineapple) Bromeliaceae	Leaf spot Root rot/heart rot ( <i>Phytophthora parasitica</i> ) Black rot	3.0L-4.0L/Ha 1.2L-2L/acre 120ml-240ml/square (Use 500L of water/Ha)	8 days	Begin spraying when weather conditions favour the development of disease, but before infection has occurred.
Tobacco	Blue Mold ( <i>Peronospora tabacina</i> )	3.0L-4.0L/Ha 1.2L-2L/acre 120ml-200ml/square (Use 500L of water/Ha)	35 days	Seed bed: 1st spray at 4- leaf stage (cross stage); last application 1-2 days before transplanting. Field: Spray 7 days after transplanting. Apply a maximum of 4 sprays per season in seed-bed and field.
Cabbage, Broccoli, Cauliflower	Damping Off ( <i>Pythium vexans</i> )	3.0L-4.0L/Ha 1.2L-2L/acre 120ml-200ml/square (Use 500L of water/Ha)	8 days	Soak seeds before sowing. Use for the first 4 sprays during the period of active growth. Protective application; 7-10 day interval in situations with heavy disease pressure.
Mustard			20 days	

PCA Registration number:

Expiration:

Batch Code:

CAUTION

CAUTION